

Motion in one dimension

21

Introduction

www ESAGS

This chapter is about how things move along a straight line or, more scientifically, how things move *in one dimension*. This is useful for learning how to describe the movement (motion) of cars along a straight road or of trains along straight railway tracks. There are three features of motion that we use to describe exactly how an object moves. They are:

Traffic often moves along a straight line.



Photograph by warrenski on Flickr.com

FACT

The **jerk** is the name we give to how fast the acceleration is changing.

1. **position** which tells us about an object's location or **displacement** which tells us about change of location
2. **speed** which tells us how fast the object is moving or **velocity** which tells us how fast the object is moving and where it is moving to, and
3. **acceleration** which tells us exactly how fast the object's speed and velocity are changing.

See introductory video: (📺 Video: VPgiq at www.everythingscience.co.za)

Reference frame

www ESAGT

The first thing to focus on when studying motion of an object or person is their position. The word *position* describes your location (where you are). However, saying that you are *here* or *there* is meaningless, you have to use known points (reference points) to help specify your position.

For example, if you were in a classroom and wanted to tell a classmate where you were standing you would first give them a reference point. The reference point might be the classroom door. You would then be able to say that you are 2 m from the doorway. This still doesn't give your position precisely. We need to provide a reference point and a coordinate system to use to define the location precisely.

Describing your location



Photograph by chokola on Flickr.com

Then you can say that you are, for example, 2 m from the door directly inside the classroom. The classroom door is a reference point and inside/outside is the coordinate system you have chosen. A frame of reference or **reference frame** is reference point which serves as the origin for a coordinate system. The coordinate system can be up or down, inside or outside, left or right or even forward or backward. These are all examples that define a 1-dimensional coordinate system. We choose one of the directions as the **positive** direction.

DEFINITION: Frame of reference

A frame of reference is a reference point combined with a set of directions.

A graphical representation of a 1-dimensional frame of reference:





Figure 21.1: Frame of reference


You can define different frames of reference for the same problem but the outcome, the physical results, will be the same. For example, a boy is standing still inside a train as it pulls out of a station. Both you and the boy define your location as the point of reference and the direction train is moving as a where you are standing as the point of reference and the direction the train is moving in as forward.

You are standing on the platform watching the train move from left to right. To you it looks as if the boy is moving from left to right, because relative to where you are standing (the platform), he is moving. According to the boy, and his *frame of reference* (the train), he is not moving.

A frame of reference must have an origin (where you are standing on the platform) and at least a positive direction. The train was moving from left to right, making to your right positive and to your left negative. If someone else was looking at the same boy, his frame of reference will be different. For example, if he was standing on the other side of the platform, the boy will be moving from right to left.



From your frame of reference the boy is moving from left to right.



For this chapter, we will only use frames of reference in the x -direction. By doing this we restrict ourselves to *one dimensional motion*. We can use the sign of the position value (positive or negative) to indicate the direction relative to the origin.

DEFINITION: *One dimensional motion*

An object is constrained to move back and forth along a line.

For example the blue dot in the figure below can only move along the x -axis.



Position




DEFINITION: Position

Position is a measurement of a location, with reference to an origin.


Quantity: Position (x) Unit name: metre Unit symbol: m

A position is a measurement of a location within a reference frame. This means that positions can be negative or positive depending on the choice for the reference frame's coordinate system. ▶ See video: VPgmf at www.everythingscience.co.za

Depending on which reference point we choose, we can say that the school is 300 m from Kosma's house (with Kosma's house as the reference point or origin) or 500 m from Kevin's house (with Kevin's house as the reference point or origin).




The shop is also 300 m from Kosma's house, but in the opposite direction as the school. When we choose a reference point, we have a positive direction and a negative direction. If we choose the direction **towards the school** as **negative**, then the direction towards the shop is positive. A negative direction is always opposite to the direction chosen as positive.



The origin is at Kosma's house and the position of the school is -300 m. Positions towards the left are defined as negative and positions towards the right are defined as positive.

Note that we could also choose the **positive** direction to be **towards the school**. In this case Kosma's house is still 300 m away from the school, but it is now in the positive direction.



The origin is at Kosma's house and the position of the school is +300 m. Positions towards the left are defined as positive and positions towards the right are defined as negative.

Group Discussion: Reference points

Divide into groups of 5 for this activity. On a straight line, choose a reference point. Since position can have both positive and negative values, discuss the advantages and disadvantages of choosing


1. either end of the line,
2. the middle of the line. (This reference point can also be called "the origin".)

Stand in a straight line, take turns choosing a different group member as the origin. Let the group member choose which direction along the line is positive. Everyone should then try to define their position. You don't need to measure your position very precisely but can just approximate it. It is important to understand whether your position is positive or negative for each difference origin and choice of coordinate system.


Notice that your position is different every time but that you didn't actually move. How you write an answer might be affected by the choice of a coordinate system but physical processes should never be affected.

Exercise 21 - 1


1. Write down the positions for objects at A, B, D and E. Do not forget the units.



2. Write down the positions for objects at F, G, H and J. Do not forget the units.



3. There are 5 houses on Newton Street, A, B, C, D and E. For all cases, assume that positions to the right are positive.



- Draw a frame of reference with house A as the origin and write down the positions of houses B, C, D and E.
- You live in house C. What is your position relative to house E?
- What are the positions of houses A, B and D, if house B is taken as the reference point?

A+ More practice video solutions **?** or help at www.everythingscience.co.za

(1.) 009g (2.) 009h (3.) 009i

Displacement and distance

ESAGV

Tip

The symbol Δ is read out as *delta*. Δ is a letter of the Greek alphabet and is used in Mathematics and Science to indicate a change in a certain quantity, or a final value minus an initial value. For example, Δx means change in x while Δt means change in t .

DEFINITION: Distance

Distance is the total length of the path taken in going from the initial position, \vec{x}_i , to the final position, \vec{x}_f . Distance is a scalar.


Quantity: Distance (D)

Unit name: metre

Unit symbol: m

In the simple map below you can see the path that winds because of a number of hills from a school to a nearby shop. The path is shown by a dashed line. The initial point, \vec{x}_i , is the school and the final point, \vec{x}_f , is the shop.

Distance is the length of dashed line. It is how far you have to walk along the path from the school to the shop.



DEFINITION: Displacement

Displacement is the change in an object's position. It is a vector that points from the initial position (\vec{x}_i) to the final position (\vec{x}_f).

Quantity: Displacement ($\Delta\vec{x}$) Unit name: metre Unit symbol: m


The displacement of an object is defined as its change in position (final position minus initial position). Displacement has a magnitude and direction and is therefore a vector. For example, if the initial position of a car is \vec{x}_i and it moves to a final position of \vec{x}_f , then the displacement is:

$$\Delta\vec{x} = \vec{x}_f - \vec{x}_i$$

To help visualise what the displacement vector looks like think back to the tail-to-head method. The displacement is the vector you add to the initial position vector to get a vector to the final position. However, subtracting an initial quantity from a final quantity happens often in Physics, so we use the shortcut Δ to mean *final - initial*. Therefore, displacement can be written:

$$\Delta\vec{x} = \vec{x}_f - \vec{x}_i$$

The following diagram illustrates the concept of displacement:



For example, if you roll a ball 5 m along a floor, in a straight line, then its displacement is 5 m, taking the direction of motion as positive, and the initial position as 0 m.

Displacement does not depend on the path travelled, but only on the initial and final positions. We use the word *distance* to describe how far an object travels along a particular path.

Tip

The words *initial* and *final* will be used very often in Physics. *Initial* refers to the situation in the beginning of the description/problem and *final* to the situation at the end. It will often happen that the final value is smaller than the initial value, such that the difference is negative. This is ok!

Tip

We will use D in this book, but you may see d used in other books.

Tip

We use the expression '*as the crow flies*' to mean a straight line between two points because birds can fly directly over many obstacles.

If we go back to the simple map repeated below you can see the path as before shown by a dashed line.

Distance is the length of dashed line. The displacement is different. Displacement is the straight-line distance from the starting point to the endpoint – from the school to the shop in the figure as shown by the solid arrow.

▶ See video: VPgmo at www.everythingscience.co.za





Illustration of distance and displacement

If we use the same situation as earlier we can explore the concepts in more detail. Consider our description of the location of the houses, school and the shop.



Komal walks to meet Kevin at his house before walking to school. What is Komal's displacement and what distance did he cover if he walks to school via Kevin's house?

Komal covers a distance of 400 m to Kevin's house and another 500 m from Kevin's house to the school. He covers a total distance of 900 m. His displacement, however, is only 100 m towards the school. This is because displacement only looks at the starting position (his house) and the end position (the school). It does not depend on the path he travelled.

To calculate his distance and displacement, we need to choose a reference point and a direction. Let's choose Komal's house as the reference point, and towards Kevin's house as the positive direction (which means that towards the school is negative). We would do the calculations as follows:

$$\begin{aligned}
 \text{Distance (D)} &= \text{path travelled} & \text{Displacement}(\Delta \vec{x}) &= \vec{x}_f - \vec{x}_i \\
 &= 400 \text{ m} + 500 \text{ m} & &= -100 \text{ m} + 0 \text{ m} \\
 &= 900 \text{ m} & &= -100 \text{ m} \\
 & & &= 100 \text{ m (in the negative } x \text{ direction)}
 \end{aligned}$$

Very often in calculations you will get a negative answer. For example, Komal's displacement in the example above, is calculated as -100 m . The minus sign in front of the answer means that his displacement is 100 m in the opposite direction (opposite to the direction chosen as positive in the beginning of the question). When we start a calculation we choose a frame of reference and a positive direction. In the first example above, the reference point is Komal's house and the positive direction is towards Kevin's house. Therefore Komal's displacement is 100 m towards the school. Notice that distance has no direction,

but displacement has a direction.

Kevin walks to school with Komal and after school walks back home. What is Kevin's displacement and what distance did he cover? For this calculation we use Kevin's house as the reference point. Let's take towards the school as the positive direction.

Distance (D) = path travelled	Displacement($\Delta\vec{x}$) = $\vec{x}_f - \vec{x}_i$
= 500 m + 500 m	= 0 m + 0 m
= 1000 m	= 0 m

It is possible to have a displacement of 0 m and a distance that is not 0 m. This happens whenever you end at the same point you started.


Differences between distance and displacement

The differences between distance and displacement can be summarised as:

Distance	Displacement
1. depends on the path	1. independent of path taken
2. always positive	2. can be positive or negative
3. is a scalar	3. is a vector

Exercise 21 - 2

1. Use this figure to answer the following questions.



- Kogis walks to Kosma's house and then to school, what is her distance and displacement?
 - Kholo walks to Kosma's house and then to school, what is her distance and displacement?
 - Komal walks to the shop and then to school, what is his distance and displacement?
 - What reference point did you use for each of the above questions?
2. You stand at the front door of your house (displacement, $\Delta\vec{x} = 0$ m). The street is 10 m away from the front door. You walk to the street and back again.

- a. What is the distance you have walked?
- b. What is your final displacement?
- c. Is displacement a vector or a scalar? Give a reason for your answer.

 More practice
  video solutions
  or help at www.everythingscience.co.za

(1.) 009j (2.) 009k

Speed and velocity

 ESAGW

DEFINITION: Average speed

Average speed is the **distance** (D) travelled divided by the time (Δt) taken for the journey.

Quantity: average speed (v_{av})

Unit name: metre per second

Unit

symbol: $\text{m} \cdot \text{s}^{-1}$

DEFINITION: Average velocity


Average velocity is the **change in position** of a body divided by the time it took for the displacement to occur.

Quantity: average velocity (\vec{v}_{av})

Unit name: metre per second

Unit

symbol: $\text{m} \cdot \text{s}^{-1}$

Before moving on review the difference between distance and displacement. Sometimes the average speed can be a very big number while the average velocity is zero.  See video:

at www.everythingscience.co.za

Average velocity is the rate of change of position. It tells us how much an object's position changes per unit of time. Velocity is a vector. We use the symbol \vec{v}_{av} for average velocity.

If we have a displacement of $\Delta\vec{x}$ and a time taken of Δt , \vec{v}_{av} is then defined as:

$$\begin{aligned}\text{average velocity (in m} \cdot \text{s}^{-1}\text{)} &= \frac{\text{change in position (in m)}}{\text{change in time (in s)}} \\ \vec{v}_{av} &= \frac{\Delta\vec{x}}{\Delta t}\end{aligned}$$

Velocity can be positive or negative. A positive velocity points in the direction you chose as positive in your coordinate system. A negative velocity points in the direction opposite to the positive direction.


Average speed (symbol v_{av}) is the distance travelled (D) divided by the time taken (Δt) for the journey. Distance and time are scalars and therefore speed will also be a scalar. Speed is calculated as follows:

$$\begin{aligned}\text{average speed (in m} \cdot \text{s}^{-1}\text{)} &= \frac{\text{distance (in m)}}{\text{time (in s)}} \\ v_{av} &= \frac{D}{\Delta t}\end{aligned}$$

Example 1: Average speed and average velocity

QUESTION

James walks 2 km away from home in 30 minutes. He then turns around and walks back home along the same path, also in 30 minutes. Calculate James' average speed and average velocity.



SOLUTION

Step 1 : Identify what information is given and what is asked for
The question explicitly gives

- the distance and time out (2 km in 30 minutes)
- the distance and time back (2 km in 30 minutes)

Step 2 : Check that all units are SI units.

The information is not in SI units and must therefore be converted.

To convert km to m, we know that:

$$1 \text{ km} = 1\,000 \text{ m}$$

$$\therefore 2 \text{ km} = 2\,000 \text{ m} \quad (\text{multiply both sides by } 2.)$$

Similarly, to convert 30 minutes to seconds,

$$1 \text{ min} = 60 \text{ s}$$

$$\therefore 30 \text{ min} = 1\,800 \text{ s} \quad (\text{multiply both sides by } 30)$$

Step 3 : Determine James' displacement and distance.

James started at home and returned home, so his displacement is 0 m.

$$\Delta \vec{x} = 0 \text{ m}$$

James walked a total distance of 4000 m (2 000 m out and 2 000 m back).

$$D = 4\,000 \text{ m}$$

Step 4 : Determine his total time.

James took 1 800 s to walk out and 1 800 s to walk back.

$$\Delta t = 3\,600 \text{ s}$$

Step 5 : Determine his average speed

$$\begin{aligned} v_{av} &= \frac{D}{\Delta t} \\ &= \frac{4\,000 \text{ m}}{3\,600 \text{ s}} \\ &= 1,11 \text{ m} \cdot \text{s}^{-1} \end{aligned}$$

Step 6 : Determine his average velocity

$$\begin{aligned} \vec{v}_{av} &= \frac{\Delta \vec{x}}{\Delta t} \\ &= \frac{0 \text{ m}}{3\,600 \text{ s}} \\ &= 0 \text{ m} \cdot \text{s}^{-1} \end{aligned}$$

Differences between speed and velocity ESAGX


The differences between speed and velocity can be summarised as:

Speed	Velocity
1. depends on the path taken	1. independent of path taken
2. always positive	2. can be positive or negative
3. is a scalar	3. is a vector
4. no dependence on direction and so is only positive	4. direction can be determined from the sign convention used (i.e. positive or negative)

Additionally, an object that makes a round trip, i.e. travels away from its starting point and then returns to the same point has zero velocity but travels at a non-zero speed.


Exercise 21 - 3

1. Bongani has to walk to the shop to buy some milk. After walking 100 m, he realises that he does not have enough money, and goes back home. If it took him two minutes to leave and come back, calculate the following:
 - a. How long was he out of the house (the time interval Δt in seconds)?
 - b. How far did he walk (distance (D))?
 - c. What was his displacement ($\Delta \vec{x}$)?
 - d. What was his average velocity (in $\text{m} \cdot \text{s}^{-1}$)?
 - e. What was his average speed (in $\text{m} \cdot \text{s}^{-1}$)?




2. Bridget is watching a straight stretch of road from her classroom window. She can see two poles which she earlier measured to be 50 m apart. Using her stopwatch, Bridget notices that it takes 3 s for most cars to travel from the one pole to the other.

- Using the equation for velocity ($\vec{v}_{av} = \frac{\Delta \vec{x}}{\Delta t}$), show all the working needed to calculate the velocity of a car travelling from the left to the right.
- If Bridget measures the velocity of a red Golf to be $-16,67 \text{ m} \cdot \text{s}^{-1}$, in which direction was the Golf travelling? Bridget leaves her stopwatch running, and notices that at $t = 5,0 \text{ s}$, a taxi passes the left pole at the same time as a bus passes the right pole. At time $t = 7,5 \text{ s}$ the taxi passes the right pole. At time $t = 9,0 \text{ s}$, the bus passes the left pole.
- How long did it take the taxi and the bus to travel the distance between the poles? (Calculate the time interval (Δt) for both the taxi and the bus).
- What was the average velocity of the taxi and the bus?
- What was the average speed of the taxi and the bus?
- What was the average speed of taxi and the bus in $\text{km} \cdot \text{h}^{-1}$?



- A rabbit runs across a freeway. There is a car, 100 m away travelling towards the rabbit.



- If the car is travelling at $120 \text{ km} \cdot \text{h}^{-1}$, what is the car's speed in $\text{m} \cdot \text{s}^{-1}$.
- How long will it take the a car to travel 100 m?
- If the rabbit is running at $10 \text{ km} \cdot \text{h}^{-1}$, what is its speed in $\text{m} \cdot \text{s}^{-1}$?

- d. If the freeway has 3 lanes, and each lane is 3 m wide, how long will it take for the rabbit to cross all three lanes?
- e. If the car is travelling in the furthestmost lane from the rabbit, will the rabbit be able to cross all 3 lanes of the freeway safely?

 More practice  video solutions  or help at www.everythingscience.co.za

(1.) 009m (2.) 009n (3.) 009p

Investigation: An exercise in safety

Divide into groups of 4 and perform the following investigation. Each group will be performing the same investigation, but the aim for each group will be different.

1. Choose an aim for your investigation from the following list and formulate a hypothesis:
 - Do cars travel at the correct speed limit?
 - Is it safe to cross the road outside of a pedestrian crossing?
 - Does the colour of your car determine the speed you are travelling at?
 - Any other relevant question that you would like to investigate.
2. On a road that you often cross, measure out 50 m along a straight section, far away from traffic lights or intersections.
3. Use a stopwatch to record the time each of 20 cars take to travel the 50 m section you measured.
4. Design a table to represent your results. Use the results to answer the question posed in the aim of the investigation. You might need to do some more measurements for your investigation. Plan in your group what else needs to be done.
5. Complete any additional measurements and write up your investigation under the following headings:
 - Aim and Hypothesis
 - Apparatus
 - Method
 - Results
 - Discussion

- Conclusion

6. Answer the following questions:

- How many cars took less than 3 s to travel 50 m?
- What was the shortest time a car took to travel 50 m?
- What was the average time taken by the 20 cars?
- What was the average speed of the 20 cars?
- Convert the average speed to $\text{km} \cdot \text{h}^{-1}$.



Acceleration



DEFINITION: Average acceleration

Average acceleration is the change in average velocity divided by the time taken.

Quantity: Average acceleration (\vec{a}_{av})
squared Unit symbol: $\text{m} \cdot \text{s}^{-2}$

Unit name: metre per second

Acceleration is a measure of how fast the velocity of an object changes in time. If we have a change in velocity ($\Delta \vec{v}$) over a time interval (Δt), then the average acceleration (\vec{a}_{av}) is defined as:

$$\text{average acceleration (in } \text{m} \cdot \text{s}^{-2}) = \frac{\text{change in velocity (in } \text{m} \cdot \text{s}^{-1})}{\text{change in time (in s)}}$$

$$\vec{a}_{av} = \frac{\Delta \vec{v}}{\Delta t}$$

We only deal with problems with constant acceleration. This means that the average acceleration and the instantaneous acceleration are the same. To make things simpler we will


only talk about acceleration and not average or instantaneous. This is represented as \vec{a} . We can also have the magnitude of the acceleration. This is:

$$a = \frac{\Delta \vec{v}}{\Delta t}$$

Acceleration is a vector. Acceleration does not provide any information about the motion, but only about how the motion changes. It is not possible to tell how fast an object is moving or in which direction from the acceleration alone.

▶ See video: VPgly at www.everythingscience.co.za

Like velocity, acceleration can be negative or positive. We see that when the sign of the acceleration and the velocity are the same, the object is speeding up. If both velocity and acceleration are positive, the object is speeding up in a positive direction. If both velocity and acceleration are negative, the object is speeding up in a negative direction. We can see this in the following diagram:



If velocity is positive and acceleration is negative, then the object is slowing down. Similarly, if the velocity is negative and the acceleration is positive the object is slowing down. This is illustrated in the following worked example.

Example 2: Acceleration

QUESTION

A car accelerates uniformly from an initial velocity of $2 \text{ m}\cdot\text{s}^{-1}$ to a final velocity of $10 \text{ m}\cdot\text{s}^{-1}$ in 8 seconds. It then slows down uniformly to a final velocity of $4 \text{ m}\cdot\text{s}^{-1}$ in 6 seconds. Calculate the acceleration of the car during the first 8 seconds and during the last 6 seconds.

SOLUTION

Tip

Avoid the use of the word *deceleration* to refer to a negative acceleration. This word usually means *slowing down* and it is possible for an object to slow down with both a positive and negative acceleration, because the sign of the velocity of the object must also be taken into account to determine whether the body is slowing down or not.

Step 1 : Choose a reference frame

We choose the point where the car starts to accelerate as the origin and the direction in which the car is already moving as the positive direction.

Step 2 : Identify what information is given and what is asked for:

Consider the motion of the car in two parts: the first 8 seconds and the last 6 seconds.

For the first 8 seconds:

$$\vec{v}_i = 2 \text{ m} \cdot \text{s}^{-1}$$

$$\vec{v}_f = 10 \text{ m} \cdot \text{s}^{-1}$$

$$t_i = 0 \text{ s}$$

$$t_f = 8 \text{ s}$$

For the last 6 seconds:

$$\vec{v}_i = 10 \text{ m} \cdot \text{s}^{-1}$$

$$\vec{v}_f = 4 \text{ m} \cdot \text{s}^{-1}$$

$$t_i = 8 \text{ s}$$

$$t_f = 14 \text{ s}$$

Step 3 : Calculate the acceleration.

For the first 8 seconds:

$$\begin{aligned} a &= \frac{\Delta v}{\Delta t} \\ &= \frac{10}{8 \text{ s} - 0 \text{ s}} \\ &= 1 \text{ m} \cdot \text{s}^{-2} \end{aligned}$$

For the next 6 seconds:

$$\begin{aligned} a &= \frac{\Delta v}{\Delta t} \\ &= \frac{4}{14 \text{ s} - 8 \text{ s}} \\ &= -1 \text{ m} \cdot \text{s}^{-2} \end{aligned}$$

During the first 8 seconds the car had a positive acceleration. This means that its velocity increased. The velocity is positive so the car is speeding up. During the next 6 seconds the car had a negative acceleration. This means that its velocity decreased. The velocity is positive so the car is slowing down.

Exercise 21 - 4

1. An athlete is accelerating uniformly from an initial velocity of $0 \text{ m} \cdot \text{s}^{-1}$ to a final velocity of $4 \text{ m} \cdot \text{s}^{-1}$ in 2 seconds. Calculate his acceleration. Let the direction that the athlete is running in be the positive direction.
2. A bus accelerates uniformly from an initial velocity of $15 \text{ m} \cdot \text{s}^{-1}$ to a final velocity of $7 \text{ m} \cdot \text{s}^{-1}$ in 4 seconds. Calculate the acceleration of the bus. Let the direction of motion of the bus be the positive direction.
3. An aeroplane accelerates uniformly from an initial velocity of $100 \text{ m} \cdot \text{s}^{-1}$ to a velocity of $200 \text{ m} \cdot \text{s}^{-1}$ in 10 seconds. It then accelerates uniformly to a final velocity of $240 \text{ m} \cdot \text{s}^{-1}$ in 20 seconds. Let the direction of motion of the aeroplane be the positive direction.
 - a. Calculate the acceleration of the aeroplane during the first 10 seconds

of the motion.

- b. Calculate the acceleration of the aeroplane during the next 20 seconds of its motion.

A+ More practice  video solutions  or help at www.everythingscience.co.za

(1.) 009q (2.) 009r (3.) 009s

Instantaneous velocity and speed

 ESAGZ

Sprinters taking off



Sprinters finishing race



Photographs by wwarby on Flickr.com

We have looked at the average velocity and speed but sometimes we want to be more precise about what is happening between the initial and final times in a problem.

Instantaneous velocity is the velocity at a specific instant in time. This can be different to the average velocity if the velocity isn't constant.

Look at the photographs of the sprinters in a race. Their velocity is different as they take off and as they end. Their average velocity for the race doesn't change but their instantaneous velocity, as captured in the "snapshots" of an instant in time does change. Their velocity in the photograph would be an instantaneous velocity.

Tip

An instant in time is different from the time taken or the time interval. It is therefore useful to use the symbol t for an instant in time (for example during the 4th second) and the symbol Δt for the time taken (for example during the first 5 seconds of the motion).

DEFINITION: Instantaneous velocity

Instantaneous velocity is the change in position over the change in a very small time interval ($\Delta t \approx 0$).

Quantity: Instantaneous velocity (\vec{v})

Unit name: metre per second

Unit name: $\text{m} \cdot \text{s}^{-1}$

DEFINITION: *Instantaneous speed*

Instantaneous speed is the magnitude of instantaneous velocity.

Quantity: Instantaneous speed (v)

Unit name: metre per second

Unit symbol: $\text{m} \cdot \text{s}^{-1}$

Instantaneous velocity is a vector. Instantaneous speed is the magnitude of instantaneous velocity. It has the same value but is not a vector so it has no direction.

Description of motion



The purpose of this chapter is to describe motion, and now that we understand the definitions of displacement, distance, velocity, speed and acceleration, we are ready to start using these ideas to describe how an object or person is moving. We will look at three ways of describing motion:

1. words
2. diagrams
3. graphs

These methods will be described in this section.

We will consider three types of motion: when the object is not moving (stationary object), when the object is moving at a constant velocity (uniform motion) and when the object is moving at a constant acceleration (motion at constant acceleration).

Stationary Object



The simplest motion that we can come across is that of a stationary object. A stationary object does not move and so its position does not change.

Vivian stands at a stop sign.

Consider an example, Vivian is waiting for a taxi. She is standing two metres from a stop street at $t = 0$ s. After one minute, at $t = 60$ s, she is still 2 metres from the stop street and after two minutes, at $t = 120$ s, also 2 metres from the stop street. Her position has not changed. Her displacement is zero (because her position is the same), her velocity is zero (because her displacement is zero) and her acceleration is also zero (because her velocity is not changing).

We can now draw graphs of position vs. time (\vec{x} vs. t), velocity vs. time (\vec{v} vs. t) and acceleration vs. time (\vec{a} vs. t) for a stationary object. The graphs are shown below.



Photograph by Rob Boudon on Flickr.com




Figure 21.2: Graphs for a stationary object (a) position vs. time (b) velocity vs. time (c) acceleration vs. time.

Vivian's position is 2 metres in the positive direction from the stop street. If the stop street is taken as the reference point, her position remains at 2 metres for 120 seconds. The graph is a horizontal line at 2 m. The velocity and acceleration graphs are also shown. They are both horizontal lines on the x -axis. Since her position is not changing, her velocity is $0 \text{ m} \cdot \text{s}^{-1}$ and since velocity is not changing, acceleration is $0 \text{ m} \cdot \text{s}^{-2}$.

DEFINITION: Gradient

(Recall from Mathematics) The gradient, m , of a line can be calculated by dividing the change in the y -value (dependent variable) by the change in the x -value (independent variable). $m = \frac{\Delta y}{\Delta x}$

Tip

The gradient of a position vs. time graph gives the average velocity, while the tangent of a position vs. time graph gives the instantaneous velocity.

Tip

The gradient of a velocity vs. time graph gives the average acceleration, while the tangent of a velocity vs. time graph gives the instantaneous acceleration.

Tip

The area under the velocity vs. time graph gives the displacement.

Since we know that velocity is the rate of change of position, we can confirm the value for the velocity vs. time graph, by calculating the gradient of the \vec{x} vs. t graph.

If we calculate the gradient of the \vec{x} vs. t graph for a stationary object we get:

$$\begin{aligned} v &= \frac{\Delta \vec{x}}{\Delta t} \\ &= \frac{\vec{x}_f - \vec{x}_i}{t_f - t_i} \\ &= \frac{2 \text{ m} - 2 \text{ m}}{120 \text{ s} - 60 \text{ s}} \text{ (initial position = final position)} \\ &= 0 \text{ m} \cdot \text{s}^{-1} \text{ (for the time that Vivian is stationary)} \end{aligned}$$

Similarly, we can confirm the value of the acceleration by calculating the gradient of the velocity vs. time graph.

If we calculate the gradient of the \vec{v} vs. t graph for a stationary object we get:

$$\begin{aligned} a &= \frac{\Delta v}{\Delta t} \\ &= \frac{\vec{v}_f - \vec{v}_i}{t_f - t_i} \\ &= \frac{0 \text{ m} \cdot \text{s}^{-1} - 0 \text{ m} \cdot \text{s}^{-1}}{120 \text{ s} - 60 \text{ s}} \\ &= 0 \text{ m} \cdot \text{s}^{-2} \end{aligned}$$

Additionally, because the velocity vs. time graph is related to the position vs. time graph, we can use the area under the velocity vs. time graph to calculate the displacement of an object.

The displacement of the object is given by the area under the graph, which is 0 m. This is obvious, because the object is not moving.

Motion at Constant Velocity




Motion at a constant velocity or *uniform motion* means that the position of the object is changing at the same rate.

Assume that Vivian takes 100 s to walk the 100 m to the taxi-stop every morning. If we assume that Vivian's house is the origin and the direction to the taxi is positive, then


Vivian's velocity is:

$$\begin{aligned}
 v &= \frac{\Delta \vec{x}}{\Delta t} \\
 &= \frac{x_f - x_i}{t_f - t_i} \\
 &= \frac{100 \text{ m} - 0 \text{ m}}{100 \text{ s} - 0 \text{ s}} \\
 &= 1 \text{ m} \cdot \text{s}^{-1}
 \end{aligned}$$

Vivian's velocity is $1 \text{ m} \cdot \text{s}^{-1}$. This means that she walked 1 m in the first second, another metre in the second second, and another in the third second, and so on. For example, after 50 s she will be 50 m from home. Her position increases by 1 m every 1 s. A diagram of Vivian's position is shown below:




We can now draw graphs of position vs. time (\vec{x} vs. t), velocity vs. time (\vec{v} vs. t) and acceleration vs. time (\vec{a} vs. t) for Vivian moving at a constant velocity. The graphs are shown here:



Graphs for motion at constant velocity (a) position vs. time (b) velocity vs. time (c) acceleration vs. time. The area of the shaded portion in the v vs. t graph corresponds to the object's displacement.

In the evening Vivian walks 100 m from the bus stop to her house in 100 s. Assume that Vivian's house is the origin. The following graphs can be drawn to describe the motion.



Graphs for motion with a constant negative velocity. The area of the shaded portion in the v vs. t graph corresponds to the object's displacement.

We see that the \vec{v} vs. t graph is a horizontal line. If the velocity vs. time graph is a horizontal line, it means that the velocity is *constant* (not changing). Motion at a constant velocity is known as *uniform motion*. We can use the \vec{x} vs. t to calculate the velocity by finding the gradient of the line.

$$\begin{aligned}
 v &= \frac{\Delta \vec{x}}{\Delta t} \\
 &= \frac{\vec{x}_f - \vec{x}_i}{t_f - t_i} \\
 &= \frac{0 \text{ m} - 100 \text{ m}}{100 \text{ s} - 0 \text{ s}} \\
 &= -1 \text{ m} \cdot \text{s}^{-1}
 \end{aligned}$$

Vivian has a velocity of $-1 \text{ m} \cdot \text{s}^{-1}$, or $1 \text{ m} \cdot \text{s}^{-1}$ towards her house. You will notice that the \vec{v} vs. t graph is a horizontal line corresponding to a velocity of $-1 \text{ m} \cdot \text{s}^{-1}$. The horizontal line means that the velocity stays the same (remains constant) during the motion. This is uniform velocity.

We can use the \vec{v} vs. t to calculate the acceleration by finding the gradient of the line.

$$\begin{aligned}
 a &= \frac{\Delta \vec{v}}{\Delta t} \\
 &= \frac{\vec{v}_f - \vec{v}_i}{t_f - t_i} \\
 &= \frac{1 \text{ m} \cdot \text{s}^{-1} - 1 \text{ m} \cdot \text{s}^{-1}}{100 \text{ s} - 0 \text{ s}} \\
 &= 0 \text{ m} \cdot \text{s}^{-2}
 \end{aligned}$$

Vivian has an acceleration of $0 \text{ m} \cdot \text{s}^{-2}$. You will notice that the graph of \vec{a} vs. t is a horizontal line corresponding to an acceleration value of $0 \text{ m} \cdot \text{s}^{-2}$. There is no acceleration during the motion because his velocity does not change.

We can use the \vec{v} vs. t graph to calculate the displacement by finding the area under the graph.

$$\begin{aligned}
 \Delta \vec{x} &= \text{Area under graph} \\
 &= \ell \times b \\
 &= 100 \text{ } (-1) \\
 &= -100 \text{ m}
 \end{aligned}$$

This means that Vivian has a displacement of 100 m towards her house.

Exercise 21 - 5

1. Use the graphs in Figure 21.6 to calculate each of the following:
 - a. Calculate Vivian's velocity between 50 s and 100 s using the x vs. t graph. Hint: Find the gradient of the line.
 - b. Calculate Vivian's acceleration during the whole motion using the v vs. t graph.
 - c. Calculate Vivian's displacement during the whole motion using the v vs. t graph.
2. Thandi takes 200 s to walk 100 m to the bus stop every morning. In the evening Thandi takes 200 s to walk 100 m from the bus stop to her home.
 - a. Draw a graph of Thandi's position as a function of time for the morning (assuming that Thandi's home is the reference point). Use the gradient of the x vs. t graph to draw the graph of velocity vs. time. Use the gradient of the v vs. t graph to draw the graph of acceleration vs. time.
 - b. Draw a graph of Thandi's position as a function of time for the evening (assuming that Thandi's home is the origin). Use the gradient of the x vs. t graph to draw the graph of velocity vs. time. Use the gradient of the v vs. t graph to draw the graph of acceleration vs. time.
 - c. Discuss the differences between the two sets of graphs in questions 2 and 3.

 More practice
  video solutions
  or help at www.everythingscience.co.za

(1.) 009t (2.) 009u

General experiment: Motion at constant velocity

Aim:

To measure the position and time during motion at constant velocity and determine the average velocity as the gradient of a "Position vs. Time" graph.

Apparatus:

A battery operated toy car, stopwatch, meter stick or measuring tape.

Method:

1. Work with a friend. Copy the table below into your workbook.
2. Complete the table by timing the car as it travels each distance.
3. Time the car twice for each distance and take the average value as your accepted time.
4. Use the distance and average time values to plot a graph of "Distance vs. Time" *onto graph paper*. Stick the graph paper into your workbook. (Remember that "A vs. B" always means "y vs. x").
5. Insert all axis labels and units onto your graph.
6. Draw the best straight line through your data points.
7. Find the gradient of the straight line. This is the average velocity.

Results:

Distance (m)	Time (s)		
	1	2	Ave.
0			
0,5			
1,0			
1,5			
2,0			
2,5			
3,0			

Conclusions:

Answer the following questions in your workbook:

1. Did the car travel with a constant velocity?


2. How can you tell by looking at the “Distance vs. Time” graph if the velocity is constant?
3. How would the “Distance vs. Time” graph look for a car with a faster velocity?
4. How would the “Distance vs. Time” graph look for a car with a slower velocity?

Motion at constant acceleration



The final situation we will be studying is motion at *constant acceleration*. We know that acceleration is the rate of change of velocity. So, if we have a constant acceleration, this means that the velocity changes at a constant rate.

Let's look at our first example of Vivian waiting at the taxi stop again. A taxi arrived and Vivian got in. The taxi stopped at the stop street and then accelerated in the positive direction as follows: After 1 s the taxi covered a distance of 2,5 m, after 2 s it covered 10 m, after 3 s it covered 22,5 m and after 4 s it covered 40 m. The taxi is covering a larger distance every second. This means that it is accelerating.



To calculate the velocity of the taxi you need to calculate the gradient of the line at each second:

$$\begin{aligned}
 \vec{v}_{1s} &= \frac{\Delta \vec{x}}{\Delta t} \\
 &= \frac{\vec{x}_f - \vec{x}_i}{t_f - t_i} \\
 &= \frac{5 \text{ m} - 0 \text{ m}}{1,5 \text{ s} - 0,5 \text{ s}} \\
 &= 5 \text{ m} \cdot \text{s}^{-1}
 \end{aligned}$$

$$\begin{aligned}
 \vec{v}_{2s} &= \frac{\Delta \vec{x}}{\Delta t} \\
 &= \frac{\vec{x}_f - \vec{x}_i}{t_f - t_i} \\
 &= \frac{15 \text{ m} - 5 \text{ m}}{2,5 \text{ s} - 1,5 \text{ s}} \\
 &= 10 \text{ m} \cdot \text{s}^{-1}
 \end{aligned}$$

$$\begin{aligned}
 \vec{v}_{3s} &= \frac{\Delta \vec{x}}{\Delta t} \\
 &= \frac{\vec{x}_f - \vec{x}_i}{t_f - t_i} \\
 &= \frac{30 \text{ m} - 15 \text{ m}}{3,5 \text{ s} - 2,5 \text{ s}} \\
 &= 15 \text{ m} \cdot \text{s}^{-1}
 \end{aligned}$$


From these velocities, we can draw the velocity-time graph which forms a straight line.

The acceleration is the gradient of the v vs. t graph and can be calculated as follows:

$$\begin{aligned}
 a &= \frac{\Delta \vec{v}}{\Delta t} \\
 &= \frac{\vec{v}_f - \vec{v}_i}{t_f - t_i} \\
 &= \frac{15 \text{ m} \cdot \text{s}^{-1} - 5 \text{ m} \cdot \text{s}^{-1}}{3 \text{ s} - 1 \text{ s}} \\
 &= 5 \text{ m} \cdot \text{s}^{-2}
 \end{aligned}$$

The acceleration does not change during the motion (the gradient stays constant). This is motion at constant or uniform acceleration.

The graphs for this situation are shown below:



Graphs for motion with a constant acceleration starting from rest.

Velocity from acceleration vs. time graphs

Just as we used velocity vs. time graphs to find displacement, we can use acceleration vs. time graphs to find the velocity of an object at a given moment in time. We simply calculate the area under the acceleration vs. time graph, at a given time. In the graph below, showing an object at a constant positive acceleration, the increase in velocity of the object after 2 seconds corresponds to the shaded portion.

$$\begin{aligned}
 v &= \text{area of rectangle} = a \times \Delta t \\
 &= 5 \text{ m} \cdot \text{s}^{-2} \times 2 \text{ s} \\
 &= 10 \text{ m} \cdot \text{s}^{-1}
 \end{aligned}$$

The velocity of the object at $t = 2 \text{ s}$ is therefore $10 \text{ m} \cdot \text{s}^{-1}$.

Summary of Graphs

The relation between graphs of position, velocity and acceleration as functions of time is summarised in the next figure.




Figure 21.3: Position-time, velocity-time and acceleration-time graphs.

Tip

The description of the motion represented by a graph should include the following (where possible):

1. whether the object is moving in the positive or negative direction
2. whether the object is at rest, moving at constant velocity or moving at constant positive acceleration (speeding up) or constant negative acceleration (slowing down)

You will also often be required to draw graphs based on a description of the motion in words or from a diagram. Remember that these are just different methods of presenting the same information. If you keep in mind the general shapes of the graphs for the different types of motion, there should not be any difficulty with explaining what is happening.


Formal experiment: *Position versus time using a ticker timer*

Aim:


To measure the position and time during motion and to use that data to plot a "Position vs. Time" graph.

Apparatus:

Trolley, ticker tape apparatus, tape, graph paper, ruler, ramp



Motion at constant velocity



Motion with increasing velocity

Method:

1. Work with a friend. Copy the table below into your workbook.
2. Attach a length of tape to the trolley.
3. Run the other end of the tape through the ticker timer.
4. Start the ticker timer going and roll the trolley down the ramp.
5. Repeat steps 1 - 3.
6. On each piece of tape, measure the distance between successive dots. Note these distances in the table below.
7. Use the frequency of the ticker timer to work out the time intervals between successive dots. Note these times in the table below.
8. Work out the average values for distance and time.
9. Use the average distance and average time values to plot a graph of "Distance vs. Time" *onto graph paper*. Stick the graph paper into your workbook. (Remember that "A vs. B" always means "y vs. x").
10. Insert all axis labels and units onto your graph.
11. Draw the best straight line through your data points.

Results:

Distance (m)			Time (s)		
1	2	Ave.	1	2	Ave.

Discussion:
Describe the motion of the trolley down the ramp.

Worked examples




The worked examples in this section demonstrate the types of questions that can be asked about graphs.

Example 3: *Description of motion based on a position-time graph*

QUESTION

The position vs. time graph for the motion of a car is given below. Draw the corresponding velocity vs. time and acceleration vs. time graphs, and then describe the motion of the car.



SOLUTION

Step 1 : Identify what information is given and what is asked for

The question gives a position vs. time graph and the following three things are required:

1. Draw a v vs. t graph.
2. Draw an a vs. t graph.
3. Describe the motion of the car.

To answer these questions, break the motion up into three sections: 0 – 2 seconds, 2 – 4 seconds and 4 – 6 seconds.

Step 2 : Velocity vs. time graph for 0 – 2 seconds

For the first 2 seconds we can see that the position (and hence the displacement) remains constant - so the object is not moving, thus it has zero velocity during this time. We can reach this conclusion by another path too: remember that the gradient of a displacement vs. time graph is the velocity. For the first 2 seconds we can see that the displacement vs. time graph is a horizontal line, i.e.. it has a gradient of zero. Thus the velocity during this time is zero and the object is stationary.

Step 3 : Velocity vs. time graph for 2 – 4 seconds


For the next 2 seconds, displacement is increasing with time so the object is moving. Looking at the gradient of the displacement graph we can see that it is not constant. In fact, the slope is getting steeper (the gradient is increasing) as time goes on. Thus, remembering that the gradient of a displacement vs. time graph is the velocity, the velocity must be increasing with time during this phase.

Step 4 : Velocity vs. time graph for 4 – 6 seconds

For the final 2 seconds we see that displacement is still increasing with time, but this time the gradient is constant, so we know that the object

is now travelling at a constant velocity, thus the velocity vs. time graph will be a horizontal line during this stage. We can now draw the graphs:

So our velocity vs. time graph looks like this one below. Because we haven't been given any values on the vertical axis of the displacement vs. time graph, we cannot figure out what the exact gradients are and therefore what the values of the velocities are. In this type of question it is just important to show whether velocities are positive or negative, increasing, decreasing or constant.



Once we have the velocity vs. time graph it's much easier to get the acceleration vs. time graph as we know that the gradient of a velocity vs. time graph is just the acceleration.

Step 5 : Acceleration vs. time graph for 0 – 2 seconds


For the first 2 seconds the velocity vs. time graph is horizontal and has a value of zero, thus it has a gradient of zero and there is no acceleration during this time. (This makes sense because we know from the displacement time graph that the object is stationary during this time, so it can't be accelerating).

Step 6 : Acceleration vs. time graph for 2 – 4 seconds

For the next 2 seconds the velocity vs. time graph has a positive gradient. This gradient is not changing (i.e. it's constant) throughout these 2 seconds so there must be a constant positive acceleration.

Step 7 : Acceleration vs. time graph for 4 – 6 seconds

For the final 2 seconds the object is travelling with a constant velocity. During this time the gradient of the velocity vs. time graph is once again zero, and thus the object is not accelerating. The acceleration vs. time graph looks like this:




Step 8 : A description of the object's motion

A brief description of the motion of the object could read something like this: At $t = 0$ s and object is stationary at some position and remains stationary until $t = 2$ s when it begins accelerating. It accelerates in a positive direction for 2 seconds until $t = 4$ s and then travels at a constant velocity for a further 2 seconds.

Example 4: Calculations from a velocity vs. time graph

QUESTION

The velocity vs. time graph of a truck is plotted below. Calculate the distance and displacement of the truck after 15 seconds.



SOLUTION

Step 1 : Decide how to tackle the problem

We are asked to calculate the distance and displacement of the car. All we need to remember here is that we can use the area between the velocity vs. time graph and the time axis to determine the distance and displacement.

Step 2 : Determine the area under the velocity vs. time graph

Break the motion up: 0 – 5 seconds, 5 – 12 seconds, 12 – 14 seconds and 14 – 15 seconds.

For 0 – 5 seconds: The displacement is equal to the area of the triangle on the left:

$$\begin{aligned}\text{Area}_{\triangle} &= \frac{1}{2} b \times h \\ &= \frac{1}{2} \times 5 \text{ s} \times 4 \text{ m} \cdot \text{s}^{-1} \\ &= 10 \text{ m}\end{aligned}$$

For 12 – 14 seconds the displacement is equal to the area of the triangle above the time axis on the right:

$$\begin{aligned}\text{Area}_{\triangle} &= \frac{1}{2} b \times h \\ &= \frac{1}{2} \times 2 \text{ s} \times 4 \text{ m} \cdot \text{s}^{-1} \\ &= 4 \text{ m}\end{aligned}$$

For 5 – 12 seconds: The displacement is equal to the area of the rectangle:

$$\begin{aligned}\text{Area}_{\square} &= \ell \times b \\ &= 7 \text{ s} \times 4 \text{ m} \cdot \text{s}^{-1} \\ &= 28 \text{ m}^2\end{aligned}$$

For 14 – 15 seconds the displacement is equal to the area of the triangle below the time axis:

$$\begin{aligned}\text{Area}_{\triangle} &= \frac{1}{2} b \times h \\ &= \frac{1}{2} \times 1 \text{ s} \times 2 \text{ m} \cdot \text{s}^{-1} \\ &= 1 \text{ m}\end{aligned}$$

Step 3 : Determine the total distance of the car

Now the total distance of the car is the sum of all of these areas:

$$\begin{aligned}D &= 10 \text{ m} + 28 \text{ m} + 4 \text{ m} + 1 \text{ m} \\ &= 43 \text{ m}\end{aligned}$$


Step 4 : Determine the total displacement of the car

Now the total displacement of the car is just the sum of all of these areas. HOWEVER, because in the last second (from $t = 14 \text{ s}$ to $t = 15 \text{ s}$) the velocity of the car is negative, it means that the car was going in the opposite direction, i.e. back where it came from! So, to find the total displacement, we have to add the first 3 areas (those with positive displacements) and subtract the last one (because it is a displacement in the opposite direction).

$$\begin{aligned}\Delta \vec{x} &= 10 \text{ m} + 28 \text{ m} + 4 \text{ m} - 1 \text{ m} \\ &= 41 \text{ m in the positive direction}\end{aligned}$$

Example 5: Velocity from a position vs. time graph
QUESTION

The position vs. time graph below describes the motion of an athlete.



1. What is the velocity of the athlete during the first 4 seconds?
2. What is the velocity of the athlete from $t = 4 \text{ s}$ to $t = 7 \text{ s}$?

SOLUTION
Step 1 : The velocity during the first 4 seconds

The velocity is given by the gradient of a position vs. time graph. During the first 4 seconds, this is


$$\begin{aligned}
 \vec{v} &= \frac{\Delta \vec{x}}{\Delta t} \\
 &= \frac{4 \text{ m} - 0 \text{ m}}{4 \text{ s} - 0 \text{ s}} \\
 &= 1 \text{ m} \cdot \text{s}^{-1}
 \end{aligned}$$

Step 2 : The velocity during the last 3 seconds

For the last 3 seconds we can see that the displacement stays constant. The graph shows a horizontal line and therefore the gradient is zero. Thus $v = 0 \text{ m} \cdot \text{s}^{-1}$.

Example 6: Drawing a v vs. t graph from an a vs. t graph
QUESTION

The acceleration vs. time graph for a car starting from rest, is given below. Calculate the velocity of the car and hence draw the velocity vs. time graph.


SOLUTION

Step 1 : Calculate the velocity values by using the area under each part of the graph.

The motion of the car can be divided into three time sections: 0 – 2 seconds; 2 – 4 seconds and 4 – 6 seconds. To be able to draw the velocity vs. time graph, the velocity for each time section needs to be calculated. The velocity is equal to the area of the square under the graph:

For 0 – 2 seconds:

$$\begin{aligned} \text{Area}_{\square} &= \ell \times b \\ &= 2 \text{ s} \times 2 \text{ m} \cdot \text{s}^{-2} \\ &= 4 \text{ m} \cdot \text{s}^{-1} \end{aligned}$$

The velocity of the car is $4 \text{ m} \cdot \text{s}^{-1}$ at $t = 2 \text{ s}$.

For 2 – 4 seconds:

$$\begin{aligned} \text{Area}_{\square} &= \ell \times b \\ &= 2 \text{ s} \times 0 \text{ m} \cdot \text{s}^{-2} \\ &= 0 \text{ m} \cdot \text{s}^{-1} \end{aligned}$$

The velocity of the car is $0 \text{ m} \cdot \text{s}^{-1}$ from $t = 2 \text{ s}$ to $t = 4 \text{ s}$.


For 4 – 6 seconds:

$$\begin{aligned} \text{Area}_{\square} &= \ell \times b \\ &= 2 \text{ s} \times -2 \text{ m} \cdot \text{s}^{-2} \\ &= -4 \text{ m} \cdot \text{s}^{-1} \end{aligned}$$

The acceleration had a negative value, which means that the velocity is decreasing. It starts at a velocity of $4 \text{ m} \cdot \text{s}^{-1}$ and decreases to $0 \text{ m} \cdot \text{s}^{-1}$.


Step 2 : Now use the values to draw the velocity vs. time graph.

The velocity vs. time graph looks like this:




Exercise 21 - 6

1. A car is parked 10 m from home for 10 minutes. Draw a displacement-time, velocity-time and acceleration-time graphs for the motion. Label all the axes.
2. A bus travels at a constant velocity of $12 \text{ m} \cdot \text{s}^{-1}$ for 6 seconds. Draw the displacement-time, velocity-time and acceleration-time graph for the motion. Label all the axes.
3. An athlete runs with a constant acceleration of $1 \text{ m} \cdot \text{s}^{-2}$ for 4 s. Draw the acceleration-time, velocity-time and displacement time graphs for the motion. Accurate values are only needed for the acceleration-time and velocity-time graphs.
4. The following velocity-time graph describes the motion of a car. Draw the displacement-time graph and the acceleration-time graph and explain the motion of the car according to the three graphs.



5. The following velocity-time graph describes the motion of a truck. Draw the displacement-time graph and the acceleration-time graph and explain the motion of the truck according to the three graphs.



More practice
 video solutions
 or help at www.everythingscience.co.za

(1.) 009v (2.) 009w (3.) 009x (4.) 009y (5.) 009z

See simulation: () Simulation: VPrwb at www.everythingscience.co.za

Equations of motion



In this section we will look at the third way to describe motion. We have looked at describing motion in terms of words and graphs. In this section we examine equations that can be used to describe motion.

This section is about solving problems relating to uniformly accelerated motion. In other words, motion at constant acceleration.

The following are the variables that will be used in this section:

$$\begin{aligned}
 \vec{v}_i &= \text{initial velocity (m} \cdot \text{s}^{-1}\text{) at } t = 0 \text{ s} \\
 \vec{v}_f &= \text{final velocity (m} \cdot \text{s}^{-1}\text{) at time } t \\
 \Delta \vec{x} &= \text{displacement (m)} \\
 t &= \text{time (s)} \\
 \Delta t &= \text{time interval (s)} \\
 \vec{a} &= \text{acceleration (m} \cdot \text{s}^{-2}\text{)}
 \end{aligned}$$

An alternate convention for some of the variables exists that you will likely encounter so

FACT

Galileo Galilei of Pisa, Italy, was the first to determine the correct mathematical law for acceleration: the total distance covered, starting from rest, is proportional to the square of the time. He also concluded that objects retain their velocity unless a force – often friction – acts upon them, refuting the accepted Aristotelian hypothesis that objects “naturally” slow down and stop unless a force acts upon them. This principle was incorporated into Newton’s laws of motion (1st law).

here is a list for reference purposes:

\vec{u} = initial velocity ($\text{m} \cdot \text{s}^{-1}$) at $t = 0 \text{ s}$

\vec{v} = final velocity ($\text{m} \cdot \text{s}^{-1}$) at time t

\vec{s} = displacement (m)

In this book we will use the first convention.

$$\vec{v}_f = \vec{v}_i + \vec{a}t \quad (21.1)$$

$$\Delta\vec{x} = \frac{(\vec{v}_i + \vec{v}_f)}{2}t \quad (21.2)$$

$$\Delta\vec{x} = \vec{v}_i t + \frac{1}{2}\vec{a}t^2 \quad (21.3)$$

$$v_f^2 = v_i^2 + 2\vec{a}\Delta\vec{x} \quad (21.4)$$

The questions can vary a lot, but the following method for answering them will always work. Use this when attempting a question that involves motion with constant acceleration. You need any three known quantities (\vec{v}_i , \vec{v}_f , $\Delta\vec{x}$, t or \vec{a}) to be able to calculate the fourth one.

Problem solving strategy:

1. Read the question carefully to identify the quantities that are given. Write them down.
2. Identify the equation to use. *Write it down!!!*
3. Ensure that all the values are in the correct units and fill them in your equation.
4. Calculate the answer and check your units.

Example 7: Equations of motion**QUESTION**

A racing car is travelling North. It accelerates uniformly covering a distance of 725 m in 10 s. If it has an initial velocity of $10 \text{ m} \cdot \text{s}^{-1}$, find its acceleration.

SOLUTION

Step 1 : **Identify what information is given and what is asked for**

We are given:

$$\begin{aligned}\vec{v}_i &= 10 \text{ m} \cdot \text{s}^{-1} \\ \Delta \vec{x} &= 725 \text{ m} \\ t &= 10 \text{ s} \\ \vec{a} &= ?\end{aligned}$$

Step 2 : Find an equation of motion relating the given information to the acceleration

If you struggle to find the correct equation, find the quantity that is not given and then look for an equation that has this quantity in it.

We can use Equation 21.3

$$\Delta \vec{x} = \vec{v}_i t + \frac{1}{2} \vec{a} t^2$$

Step 3 : Substitute your values in and find the answer

$$\begin{aligned}\Delta \vec{x} &= \vec{v}_i t + \frac{1}{2} \vec{a} t^2 \\ 725 \text{ m} &= (10 \text{ m} \cdot \text{s}^{-1} \times 10 \text{ s}) + \frac{1}{2} \vec{a} \times (10 \text{ s})^2 \\ 725 \text{ m} - 100 \text{ m} &= (50 \text{ s}^2) \vec{a} \\ \vec{a} &= 12,5 \text{ m} \cdot \text{s}^{-2}\end{aligned}$$

Step 4 : Quote the final answer

The racing car is accelerating at $12,5 \text{ m} \cdot \text{s}^{-2}$ North.

Example 8: Equations of motion I

QUESTION

A motorcycle, travelling East, starts from rest, moves in a straight line with a constant acceleration and covers a distance of 64 m in 4 s. Calculate

- its acceleration
- its final velocity
- at what time the motorcycle had covered half the total distance
- what distance the motorcycle had covered in half the total time.

SOLUTION

Step 1 : **Identify what information is given and what is asked for**

We are given:

$$\vec{v}_i = 0 \text{ m} \cdot \text{s}^{-1} \text{ (because the object starts from rest.)}$$

$$\Delta \vec{x} = 64 \text{ m}$$

$$t = 4 \text{ s}$$

$$\vec{a} = ?$$

$$\vec{v}_f = ?$$

$$t = ? \text{ at half the distance } \Delta \vec{x} = 32 \text{ m.}$$

$$\Delta \vec{x} = ? \text{ at half the time } t = 2 \text{ s.}$$

All quantities are in SI units.

Step 2 : **Acceleration: Find a suitable equation to calculate the acceleration**

We can use Equations 21.3

$$\Delta \vec{x} = \vec{v}_i t + \frac{1}{2} \vec{a} t^2$$

Step 3 : **Substitute the values and calculate the acceleration**

$$\Delta \vec{x} = \vec{v}_i t + \frac{1}{2} \vec{a} t^2$$

$$64 \text{ m} = (0 \text{ m} \cdot \text{s}^{-1} \times 4 \text{ s}) + \frac{1}{2} \vec{a} \times (4 \text{ s})^2$$

$$64 \text{ m} = (8 \text{ s}^2) \vec{a}$$

$$\vec{a} = 8 \text{ m} \cdot \text{s}^{-2} \text{ East}$$

Step 4 : **Final velocity: Find a suitable equation to calculate the final velocity**

We can use Equation 21.1 - remember we now also know the acceleration of the object.

$$\vec{v}_f = \vec{v}_i + \vec{a} t$$

Step 5 : **Substitute the values and calculate the final velocity**

$$\begin{aligned}\vec{v}_f &= \vec{v}_i + at \\ \vec{v}_f &= 0 \text{ m} \cdot \text{s}^{-1} + (8 \text{ m} \cdot \text{s}^{-2})(4 \text{ s}) \\ &= 32 \text{ m} \cdot \text{s}^{-1} \text{ East}\end{aligned}$$

Step 6 : **Time at half the distance: Find an equation to calculate the time**

We can use Equation 21.3:

$$\begin{aligned}\Delta \vec{x} &= \vec{v}_i t + \frac{1}{2} \vec{a} t^2 \\ 32 \text{ m} &= (0 \text{ m} \cdot \text{s}^{-1})t + \frac{1}{2}(8 \text{ m} \cdot \text{s}^{-2})(t)^2 \\ 32 \text{ m} &= 0 + (4 \text{ m} \cdot \text{s}^{-2})t^2 \\ 8 \text{ s}^2 &= t^2 \\ t &= 2,83 \text{ s}\end{aligned}$$

Step 7 : **Distance at half the time: Find an equation to relate the distance and time**

Half the time is 2 s, thus we have \vec{v}_i , \vec{a} and t - all in the correct units. We can use Equation 21.3 to get the distance:

$$\begin{aligned}\Delta \vec{x} &= \vec{v}_i t + \frac{1}{2} \vec{a} t^2 \\ &= (0 \text{ m} \cdot \text{s}^{-1})(2 \text{ s}) + \frac{1}{2}(8 \text{ m} \cdot \text{s}^{-2})(2 \text{ s})^2 \\ &= 16 \text{ m East}\end{aligned}$$

Exercise 21 - 7

1. A car starts off at $10 \text{ m} \cdot \text{s}^{-1}$ and accelerates at $1 \text{ m} \cdot \text{s}^{-2}$ for 10 s. What is its final velocity?

2. A train starts from rest, and accelerates at $1 \text{ m}\cdot\text{s}^{-2}$ for 10 s. How far does it move?
3. A bus is going $30 \text{ m}\cdot\text{s}^{-1}$ and stops in 5 s. What is its stopping distance for this speed?
4. A racing car going at $20 \text{ m}\cdot\text{s}^{-1}$ stops in a distance of 20 m. What is its acceleration?
5. A ball has a uniform acceleration of $4 \text{ m}\cdot\text{s}^{-1}$. Assume the ball starts from rest. Determine the velocity and displacement at the end of 10 s.
6. A motorcycle has a uniform acceleration of $4 \text{ m}\cdot\text{s}^{-1}$. Assume the motorcycle has an initial velocity of $20 \text{ m}\cdot\text{s}^{-1}$. Determine the velocity and displacement at the end of 12 s.
7. An aeroplane accelerates uniformly such that it goes from rest to $144 \text{ km}\cdot\text{hr}^{-1}$ in 8 s. Calculate the acceleration required and the total distance that it has travelled in this time.

 More practice  video solutions  or help at www.everythingscience.co.za

(1.) 00a0 (2.) 00a1 (3.) 00a2 (4.) 00a3 (5.) 00a4 (6.) 00a5
(7.) 00a6

Extension: Finding the equations of motion



The following does not form part of the syllabus and can be considered additional information.

Derivation of 21.1

According to the definition of acceleration:


$$\vec{a} = \frac{\Delta \vec{v}}{t}$$

where $\Delta \vec{v}$ is the change in velocity, i.e. $\Delta v = \vec{v}_f - \vec{v}_i$. Thus we have

$$\begin{aligned}\vec{a} &= \frac{\vec{v}_f - \vec{v}_i}{t} \\ \vec{v}_f &= \vec{v}_i + \vec{a}t\end{aligned}$$

Derivation of 21.2

We have seen that displacement can be calculated from the area under a velocity vs. time graph. For *uniformly accelerated motion* the most complicated velocity vs. time graph we can have is a straight line. Look at the graph below - it represents an object with a starting velocity of \vec{v}_i , accelerating to a final velocity \vec{v}_f over a total time t .



To calculate the final displacement we must calculate the area under the graph - this is just the area of the rectangle added to the area of the triangle. This portion of the graph has been shaded for clarity.

$$\begin{aligned}\text{Area}_{\triangle} &= \frac{1}{2}b \times h \\ &= \frac{1}{2}t \times (v_f - v_i) \\ &= \frac{1}{2}v_f t - \frac{1}{2}v_i t\end{aligned}$$

$$\begin{aligned}\text{Area}_{\square} &= \ell \times b \\ &= t \times v_i \\ &= v_i t\end{aligned}$$

$$\begin{aligned}
 \text{Displacement} &= \text{Area}_{\square} + \text{Area}_{\triangle} \\
 \Delta \vec{x} &= v_i t + \frac{1}{2} v_f t - \frac{1}{2} v_i t \\
 \Delta \vec{x} &= \frac{(v_i + v_f)}{2} t
 \end{aligned}$$

Derivation of 21.3

This equation is simply derived by eliminating the final velocity v_f in 21.2. Remembering from 21.1 that

$$\vec{v}_f = \vec{v}_i + \vec{a}t$$

then 21.2 becomes

$$\begin{aligned}
 \Delta \vec{x} &= \frac{\vec{v}_i + \vec{v}_i + \vec{a}t}{2} t \\
 &= \frac{2\vec{v}_i t + \vec{a}t^2}{2} \\
 \Delta \vec{x} &= \vec{v}_i t + \frac{1}{2} \vec{a}t^2
 \end{aligned}$$

Derivation of 21.4

This equation is just derived by eliminating the time variable in the above equation. From 21.1 we know

$$t = \frac{\vec{v}_f - \vec{v}_i}{a}$$

Substituting this into 21.3 gives

$$\begin{aligned}
 \Delta \vec{x} &= \vec{v}_i \left(\frac{\vec{v}_f - \vec{v}_i}{a} \right) + \frac{1}{2} a \left(\frac{\vec{v}_f - \vec{v}_i}{a} \right)^2 \\
 &= \frac{\vec{v}_i \vec{v}_f}{a} - \frac{\vec{v}_i^2}{a} + \frac{1}{2} a \left(\frac{\vec{v}_f^2 - 2\vec{v}_i \vec{v}_f + \vec{v}_i^2}{a^2} \right) \\
 &= \frac{\vec{v}_i \vec{v}_f}{a} - \frac{\vec{v}_i^2}{a} + \frac{\vec{v}_f^2}{2a} - \frac{\vec{v}_i \vec{v}_f}{a} + \frac{\vec{v}_i^2}{2a} \\
 2a\Delta \vec{x} &= -2\vec{v}_i^2 + \vec{v}_f^2 + \vec{v}_i^2 \\
 \vec{v}_f^2 &= \vec{v}_i^2 + 2a\Delta \vec{x}
 \end{aligned}$$

This gives us the final velocity in terms of the initial velocity, acceleration and displacement and is independent of the time variable.

Applications in the real-world



What we have learnt in this chapter can be directly applied to road safety. We can analyse the relationship between speed and stopping distance. The following worked example illustrates this application.

Example 9: Stopping distance


QUESTION

A truck is travelling at a constant velocity of $10 \text{ m} \cdot \text{s}^{-1}$ when the driver sees a child 50 m in front of him in the road. He hits the brakes to stop the truck. The truck accelerates at a rate of $-1.25 \text{ m} \cdot \text{s}^{-2}$. His reaction time to hit the brakes is 0,5 seconds. Will the truck hit the child?

SOLUTION

Step 1 : **Analyse the problem and identify what information is given**

It is useful to draw a time-line like this one:



We need to know the following:

- What distance the driver covers before hitting the brakes.
- How long it takes the truck to stop after hitting the brakes.
- What total distance the truck covers to stop.

Step 2 : Calculate the distance AB

Before the driver hits the brakes, the truck is travelling at constant velocity. There is no acceleration and therefore the equations of motion are not used. To find the distance travelled, we use:

$$v = \frac{d}{t}$$

$$10 = \frac{d}{0,5}$$

$$d = 5 \text{ m}$$

The truck covers 5 m before the driver hits the brakes.

Step 3 : Calculate the time BC

We have the following for the motion between B and C:

$$\vec{v}_i = 10 \text{ m} \cdot \text{s}^{-1}$$

$$\vec{v}_f = 0 \text{ m} \cdot \text{s}^{-1}$$

$$a = -1,25 \text{ m} \cdot \text{s}^{-2}$$

$$t = ?$$

We can use Equation 21.1

$$\vec{v}_f = \vec{v}_i + at$$

$$0 = 10 \text{ m} \cdot \text{s}^{-1} + (-1,25 \text{ m} \cdot \text{s}^{-2})t$$

$$-10 \text{ m} \cdot \text{s}^{-1} = (-1,25 \text{ m} \cdot \text{s}^{-2})t$$

$$t = 8 \text{ s}$$

Step 4 : Calculate the distance BC

For the distance we can use Equation 21.2 or Equation 21.3. We will

use Equation 21.2:

$$\begin{aligned}\Delta \vec{x} &= \frac{(\vec{v}_i + \vec{v}_f)}{2} t \\ \Delta \vec{x} &= \frac{10 + 0}{s}(8) \\ \Delta \vec{x} &= 40 \text{ m}\end{aligned}$$

Step 5 : Write the final answer

The total distance that the truck covers is $d_{AB} + d_{BC} = 5 + 40 = 45$ meters. The child is 50 meters ahead. The truck will not hit the child.

Chapter 21 | Summary

See the summary presentation (📺 Presentation: VPgjl at www.everythingscience.co.za)

- A reference point is a point from where you take your measurements.
- A frame of reference is a reference point with a set of directions.
- Your position is where you are located with respect to your reference point.
- The displacement of an object is how far it is from the reference point. It is the shortest distance between the object and the reference point. It has magnitude and direction because it is a vector.
- The distance of an object is the length of the path travelled from the starting point to the end point. It has magnitude only because it is a scalar.
- Speed (v) is the distance covered (D) divided by the time taken (Δt):

$$v = \frac{D}{\Delta t}$$

- Average velocity (\vec{v}_{av}) is the displacement ($\Delta \vec{x}$) divided by the time taken (Δt):

$$\vec{v}_{av} = \frac{\Delta \vec{x}}{\Delta t}$$

- Instantaneous speed is the speed at a specific instant in time.
- Instantaneous velocity is the velocity at a specific instant in time.

- Acceleration (\vec{a}) is the change in velocity ($\Delta\vec{v}$) over a time interval (Δt):

$$\vec{a} = \frac{\Delta\vec{v}}{\Delta t}$$

- The gradient of a position - time graph (x vs. t) gives the velocity.
- The gradient of a velocity - time graph (v vs. t) gives the acceleration.
- The area under a velocity - time graph (v vs. t) gives the displacement.
- The area under an acceleration - time graph (a vs. t) gives the velocity.
- The graphs of motion are summarised in Figure 21.3.
- The equations of motion are used where constant acceleration takes place:

$$v_f = v_i + at$$

$$\Delta\vec{x} = \frac{(v_i + v_f)}{2}t$$

$$\Delta\vec{x} = v_i t + \frac{1}{2}at^2$$

$$\vec{v}_f^2 = \vec{v}_i^2 + 2a\Delta\vec{x}$$

Physical Quantities			
Quantity	Vector	Unit name	Unit symbol
Position (x)	-	metre	m
Distance (D)	-	metre	m
Displacement ($\Delta\vec{x}$)	✓	metre	m
Speed (v_{av})	-	metre per second	$\text{m} \cdot \text{s}^{-1}$
Average velocity (\vec{v}_{av})	✓	metre per second	$\text{m} \cdot \text{s}^{-1}$
Instantaneous velocity (\vec{v})	✓	metre per second	$\text{m} \cdot \text{s}^{-1}$
Instantaneous speed (v)	-	metre per second	$\text{m} \cdot \text{s}^{-1}$
Instantaneous acceleration (\vec{a})	✓	metre per second per second	$\text{m} \cdot \text{s}^{-2}$
Average acceleration (\vec{a}_{av})	✓	metre per second per second	$\text{m} \cdot \text{s}^{-2}$
Magnitude of acceleration (a)	-	metre per second per second	$\text{m} \cdot \text{s}^{-2}$


Table 21.1: Units used in **motion in one dimension**

Chapter 21**End of chapter exercises**


1. Give one word/term for the following descriptions.
 - a. The shortest path from start to finish.
 - b. A physical quantity with magnitude and direction.
 - c. The quantity defined as a change in velocity over a time period.
 - d. The point from where you take measurements.
 - e. The distance covered in a time interval.
 - f. The velocity at a specific instant in time.
2. Choose an item from column B that match the description in column A. Write down only the letter next to the question number. You may use an item from column B more than once.

Column A	Column B
a. The area under a velocity - time graph	gradient
b. The gradient of a velocity - time graph	area
c. The area under an acceleration - time graph	velocity
d. The gradient of a position - time graph	displacement
	acceleration
	slope

3. Indicate whether the following statements are TRUE or FALSE. Write only "true" or "false". If the statement is false, write down the correct statement.
 - a. A scalar is the displacement of an object over a time interval.
 - b. The position of an object is where it is located.
 - c. The sign of the velocity of an object tells us in which direction it is travelling.
 - d. The acceleration of an object is the change of its displacement over a period in time.
4. [SC 2003/11] A body accelerates uniformly from rest for t_0 seconds after which it continues with a constant velocity. Which graph is the correct representation of the body's motion?




5. [SC 2003/11] The velocity-time graphs of two cars are represented by P and Q as shown



The difference in the distance travelled by the two cars (in m) after 4 s is ...

- (a) 12
(b) 6
(c) 2
(d) 0
6. [IEB 2005/11 HG] The graph that follows shows how the speed of an athlete varies with time as he sprints for 100 m.




Which of the following equations can be used to correctly determine the time t for which he accelerates?


- (a) $100 = (10)(11) - \frac{1}{2}(10)t$
(b) $100 = (10)(11) + \frac{1}{2}(10)t$
(c) $100 = 10t + \frac{1}{2}(10)t^2$
(d) $100 = \frac{1}{2}(0)t + \frac{1}{2}(10)t^2$
7. [SC 2002/03 HG1] In which one of the following cases will the distance covered and the magnitude of the displacement be the same?
- (a) A girl climbs a spiral staircase.
(b) An athlete completes one lap in a race.
(c) A raindrop falls in still air.

(d) A passenger in a train travels from Cape Town to Johannesburg.

8. [SC 2003/11] A car, travelling at constant velocity, passes a stationary motor cycle at a traffic light. As the car overtakes the motorcycle, the motorcycle accelerates uniformly from rest for 10 s. The following displacement-time graph represents the motions of both vehicles from the traffic light onwards.




- Use the graph to find the magnitude of the constant velocity of the car.
 - Use the information from the graph to show by means of calculation that the magnitude of the acceleration of the motorcycle, for the first 10 s of its motion is $7,5 \text{ m} \cdot \text{s}^{-2}$.
 - Calculate how long (in seconds) it will take the motorcycle to catch up with the car (point X on the time axis).
 - How far behind the motorcycle will the car be after 15 seconds?
9. [IEB 2005/11 HG] Which of the following statements is *true* of a body that accelerates uniformly?
- Its rate of change of position with time remains constant.
 - Its position changes by the same amount in equal time intervals.
 - Its velocity increases by increasing amounts in equal time intervals.
 - Its rate of change of velocity with time remains constant.
10. [IEB 2003/11 HG1] The velocity-time graph for a car moving along a straight horizontal road is shown below.



Which of the following expressions gives the magnitude of the average velocity of the car?

- (a) $\frac{\text{Area A}}{t}$
- (b) $\frac{\text{Area A} + \text{Area B}}{t}$
- (c) $\frac{\text{Area B}}{t}$
- (d) $\frac{\text{Area A} - \text{Area B}}{t}$

11. [SC 2002/11 SG] A car is driven at $25 \text{ m} \cdot \text{s}^{-1}$ in a municipal area. When the driver sees a traffic officer at a speed trap, he realises he is travelling too fast. He immediately applies the brakes of the car while still 100 m away from the speed trap.
- (a) Calculate the magnitude of the minimum acceleration which the car must have to avoid exceeding the speed limit, if the municipal speed limit is $16.6 \text{ m} \cdot \text{s}^{-1}$.
 - (b) Calculate the time from the instant the driver applied the brakes until he reaches the speed trap. Assume that the car's velocity, when reaching the trap, is $16.6 \text{ m} \cdot \text{s}^{-1}$.
12. A traffic officer is watching his speed trap equipment at the bottom of a valley. He can see cars as they enter the valley 1 km to his left until they leave the valley 1 km to his right. Nelson is recording the times of cars entering and leaving the valley for a school project. Nelson notices a white Toyota enter the valley at 11:01:30 and leave the valley at 11:02:42. Afterwards, Nelson hears that the traffic officer recorded the Toyota doing $140 \text{ km} \cdot \text{hr}^{-1}$.
- (a) What was the time interval (Δt) for the Toyota to travel through the valley?
 - (b) What was the average speed of the Toyota?
 - (c) Convert this speed to $\text{km} \cdot \text{hr}^{-1}$.
 - (d) Discuss whether the Toyota could have been travelling at $140 \text{ km} \cdot \text{hr}^{-1}$ at the bottom of the valley.
 - (e) Discuss the differences between the instantaneous speed (as measured by the speed trap) and average speed (as measured by Nelson).
13. [IEB 2003/11 HG] A velocity-time graph for a ball rolling along a track is shown below. The graph has been divided up into 3 sections, A, B and C for easy reference. (Disregard any effects of friction.)



- (a) Use the graph to determine the following:
 - i. the speed 5 s after the start
 - ii. the distance travelled in Section A
 - iii. the acceleration in Section C
 - (b) At time t_1 the velocity-time graph intersects the time axis. Use an appropriate equation of motion to calculate the value of time t_1 (in s).
 - (c) Sketch a displacement-time graph for the motion of the ball for these 12 s. (You do not need to calculate the actual values of the displacement for each time interval, but do pay attention to the general shape of this graph during each time interval.)
- 14.** In towns and cities, the speed limit is $60 \text{ km}\cdot\text{hr}^{-1}$. The length of the average car is 3.5 m, and the width of the average car is 2 m. In order to cross the road, you need to be able to walk further than the width of a car, before that car reaches you. To cross safely, you should be able to walk at least 2 m further than the width of the car (4 m in total), before the car reaches you.
- (a) If your walking speed is $4 \text{ km}\cdot\text{hr}^{-1}$, what is your walking speed in $\text{m}\cdot\text{s}^{-1}$?
 - (b) How long does it take you to walk a distance equal to the width of the average car?
 - (c) What is the speed in $\text{m}\cdot\text{s}^{-1}$ of a car travelling at the speed limit in a town?
 - (d) How many metres does a car travelling at the speed limit travel, in the same time that it takes you to walk a distance equal to the width of car?
 - (e) Why is the answer to the previous question important?
 - (f) If you see a car driving toward you, and it is 28 m away (the same as the length of 8 cars), is it safe to walk across the road?
 - (g) How far away must a car be, before you think it might be safe to cross? How many car-lengths is this distance?
- 15.** A bus on a straight road starts from rest at a bus stop and accelerates at $2 \text{ m}\cdot\text{s}^{-2}$ until it reaches a speed of $20 \text{ m}\cdot\text{s}^{-1}$. Then the bus travels for 20 s at a constant speed until the driver sees the next bus stop in the distance. The driver applies the brakes, stopping the bus in a uniform manner in 5 s.
- (a) How long does the bus take to travel from the first bus stop to the second bus stop?
 - (b) What is the average velocity of the bus during the trip?



More practice



video solutions

or help at www.everythingscience.co.za

- (1.) 00a7 (2.) 00a8 (3.) 00a9 (4.) 00aa (5.) 00ab (6.) 00ac
(7.) 00ad (8.) 00ae (9.) 00af (10.) 00ag (11.) 00ah (12.) 00ai
(13.) 00aj (14.) 00ak (15.) 00am